

Children's Vaccine Schedule

Birth through 6 years

Birth

- ☐ Hepatitis B
- ☐ RSV


2 months

- ☐ Hepatitis B
- ☐ Rotavirus
- ☐ DTaP
- ☐ Hib
- ☐ Pneumococcal
- ☐ Polio

4 months

- ☐ Rotavirus
- ☐ DTaP
- ☐ Hib
- ☐ Pneumococcal
- ☐ Polio


6 months

- ☐ Hepatitis B
- ☐ Rotavirus
- ☐ DTaP
- ☐ Hib
- ☐ Pneumococcal
- ☐ Polio


12 months

- ☐ Chickenpox
- ☐ MMR
- ☐ Hib
- ☐ Pneumococcal
- ☐ Hepatitis A

15 months

- ☐ DTaP
- ☐ Hib
- ☐ Pneumococcal

18 months

- ☐ Hepatitis A

4–6 years
Kindergarten

- ☐ DTaP
- ☐ Polio
- ☐ MMR
- ☐ Chickenpox


Yearly
Starting at 6 months

- ☐ Flu (2 doses 1 month apart when getting flu vaccine for the first time before age 9)
- ☐ COVID-19 (should get when age 6–23 months, may get when age 2+ years)



Some vaccines may be given as combo shots to reduce the number of pokes.

Talk to your healthcare provider if your child:

- Misses a dose of vaccine.
- Has a health condition that increases their risk of infection.
- Will travel outside the United States.



Tacoma-Pierce County
Health Department
Healthy People in Healthy Communities

Learn more at tpchd.org/kidsvax